

Building surrogate models of nuclear density functional theory with Gaussian processes and autoencoders - Supplementary Material

Verriere, Marc

Supplement / Prilog

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:217:924095>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-02-10**



Repository / Repozitorij:

[Repository of the Faculty of Science - University of Zagreb](#)



Building Surrogate Models of Nuclear Density Functional Theory with Gaussian Processes and Autoencoders

Supplementary Material

Marc Verriere¹, Nicolas Schunck^{1,*}, Irene Kim^{2,3}, Petar Marević^{1,4,5}, Kevin Quinlan⁶, Michelle N’Go^{6,7}, David Regnier^{8,9}, Raphael David Lasserri⁴

¹*Nuclear and Data Theory group, Nuclear and Chemical Science Division, Lawrence Livermore National Laboratory, Livermore, CA-94550, USA,*

²*Machine Intelligence group, Center for Applied Scientific Computing, Lawrence Livermore National Laboratory, Livermore, CA-94550, USA,*

³*Sensing and Intelligent Systems group, Global Security Computing Applications Division, Lawrence Livermore National Laboratory, Livermore, CA-94550, USA,*

⁴*Centre Borelli, ENS Paris-Saclay, Université Paris-Saclay, 91190 Gif-sur-Yvette, France,*

⁵*Physics Department, Faculty of Science, University of Zagreb, 10000 Zagreb, Croatia,*

⁶*Applied Statistics Group, Lawrence Livermore National Laboratory, Livermore, CA 94551, United States of America,*

⁷*Center for Complex Biological Systems, University of California Irvine, Irvine, CA 92697, USA,*

⁸*Université Paris-Saclay, CEA, Laboratoire Matière en Conditions Extrêmes, 91680 Bruyères-le-Châtel, France,*

⁹*CEA, DAM, DIF, 91297 Arpajon, France*

Correspondence*:
Corresponding Author
schunck1@llnl.gov

1 OVERLAP BETWEEN NON-ORTHOGONAL CANONICAL BASES

2 In Haider and Gogny (1992) the norm overlap between two different HFB vacua is expressed as a function
3 of the single-particle overlap between the two respective sets of canonical wavefunctions and occupation
4 numbers. This formula implicitly assumes that the canonical wavefunctions are orthonormal. When using
5 canonical wavefunctions reconstructed by the AE, this property may not hold anymore and the Haider &
6 Gogny formula should not be used ‘as is’ when evaluating the norm overlap. In this appendix, we show
7 how to generalize it.

8 We recall that for a system with conserved time-reversal symmetry, the quasiparticle vacuum can be
9 written Ring and Schuck (2004); Schunck (2019)

$$|\Phi\rangle = \prod_{\mu>} \beta_{\mu}^{\dagger} \beta_{\bar{\mu}}^{\dagger} |0\rangle, \quad (1)$$

10 where the quasiparticle operators ($\hat{\beta}, \hat{\beta}^\dagger$) are obtained from a single-particle basis (\hat{c}, \hat{c}^\dagger) by the Bogoliubov
 11 transformation W of (14) in Sec. 2.5. The Bloch-Messiah-Zumino decomposition of the matrix W is a
 12 consequence of the fact that the quasiparticle operators should obey the same anticommutation relations as
 13 the particle operators. In the resulting canonical basis, the quasiparticle vacuum can be written in the BCS
 14 form,

$$|\Phi\rangle = \prod_{\mu>} \left(u_\mu + v_\mu \hat{a}_\mu^\dagger \hat{a}_{\bar{\mu}}^\dagger \right) |0\rangle, \quad (2)$$

15 where $|0\rangle$ is the particle vacuum, $\bar{\mu}$ refers to the time-reversed partner of the state with index μ , and the
 16 summation runs only over “positive” indices μ . Implicit in this expression is the fact that the canonical
 17 wavefunctions associated with the operators (\hat{a}, \hat{a}^\dagger) are orthonormal or, equivalently, that the operators
 18 (\hat{a}, \hat{a}^\dagger) anticommute. If these conditions are not verified, the form (2) is not valid and the formulas for
 19 the norm overlap given in Haider and Gogny (1992) cannot apply. In our case, the fit of the AE gives a
 20 set of *reconstructed* canonical wavefunctions which we noted $\{\phi_\mu(\mathbf{r}, \sigma)\}_\mu$ and are associated with a set
 21 of single-particle creation and annihilation operators (\hat{f}, \hat{f}^\dagger). Although we call these objects canonical
 22 orbitals, this is somewhat a misnomer since the wavefunctions are not necessarily orthonormal. As a
 23 consequence, one cannot define a BCS state (2) with these operators. Our goal is to find a transformation
 24 of these single-particle operators that allows us to define a BCS state.

25 Following the notations of Haider and Gogny (1992), we thus define the single-particle overlaps

$$\tilde{\tau}_{\mu\nu}^{(ff)} = \{\hat{f}_\mu^\dagger, \hat{f}_\nu\} = \sum_\sigma \int d^3\mathbf{r} \phi_\mu^*(\mathbf{r}, \sigma) \phi_\nu(\mathbf{r}, \sigma). \quad (3)$$

26 The set of all such overlaps define the overlap matrix $\tilde{\tau}^{(ff)}$. This overlap matrix is block diagonal as it
 27 satisfies the relations

$$\tilde{\tau}_{\mu\nu}^{(ff)} = \tilde{\tau}_{\bar{\mu}\bar{\nu}}^{(ff)*}, \quad \tilde{\tau}_{\mu\bar{\nu}}^{(ff)} = \tilde{\tau}_{\bar{\mu}\nu}^{(ff)} = 0. \quad (4)$$

From the s.p. operators (\hat{f}, \hat{f}^\dagger), we can introduce a new set of q.p. operators ($\hat{\chi}, \hat{\chi}^\dagger$) through

$$\hat{\chi}_\mu = \tilde{u}_\mu \hat{f}_\mu - \tilde{v}_\mu \hat{f}_{\bar{\mu}}^\dagger, \quad (5a)$$

$$\hat{\chi}_{\bar{\mu}} = \tilde{u}_\mu \hat{f}_{\bar{\mu}} + \tilde{v}_\mu \hat{f}_\mu^\dagger. \quad (5b)$$

28 It is easy to see that these q.p. operators do not obey the Fermion anticommutation relation. In other words,
 29 the q.p. *spinors* associated with these operators are not orthogonal. We thus introduce the overlap matrix
 30 $\tilde{\tau}_{\mu\nu}^{(\chi\chi)}$ between any two such spinors $\mu, \nu > 0$. Owing to (5a)-(5b) and (3) it is straightforward to show that
 31 it is given by

$$\tilde{\tau}_{\mu\nu}^{(\chi\chi)} = \tilde{u}_\mu \tilde{u}_\nu \tilde{\tau}_{\mu\nu}^{(ff)} + \tilde{v}_\mu \tilde{v}_\nu \tilde{\tau}_{\mu\nu}^{(ff)*}, \quad (6)$$

32 and verify the same properties (4) as the single-particle overlap. We symmetrically orthogonalize the q.p.
 33 basis by eigendecomposing $\tilde{\tau}_{\mu\nu}^{(\chi\chi)}$ ¹

$$\tilde{\tau}_{\mu\nu}^{(\chi\chi)} = Q \Sigma^2 Q^\dagger, \quad (7)$$

¹ In fact, we can limit ourself to compute the Cholesky decomposition of $\tilde{\tau}_{\mu\nu}^{(\chi\chi)}$, but we use the eigenvalues to check the rank and invert the matrix.

34 where Q is unitary and $\Sigma = \text{diag}(\sigma_0, \sigma_1, \dots)$ with $\sigma_\mu > 0$. We then construct a new orthogonal q.p. basis

$$\hat{\gamma}_\mu^\dagger = \sum_k \hat{\chi}_k^\dagger (Q\Sigma^{-1})_{k\mu} \quad (8)$$

35 that satisfies the fermion commutation relations, $\{\hat{\gamma}_\mu^\dagger, \hat{\gamma}_\nu\} = \delta_{\mu\nu}$. We can associate with these new q.p.
36 operators $(\hat{\gamma}_\mu^\dagger, \hat{\gamma}_\nu)$ a quasiparticle vacuum of the type (1). We now need to find the Bogoliubov transformation
37 \tilde{W} (and its Bloch-Messiah decomposition) that relate the $(\hat{\gamma}_\mu^\dagger, \hat{\gamma}_\nu)$ to a properly orthonormal s.p. basis. To
38 this end, we first diagonalize the single-particle overlap matrix

$$\tilde{\tau}_{\mu\nu}^{(ff)} = R\tilde{\Sigma}^2 R^\dagger, \quad (9)$$

which defines a new set of *particle* operators $(\hat{b}, \hat{b}^\dagger)$ through the relations

$$\hat{b}_i^\dagger = \sum_k \hat{f}_k^\dagger (R\tilde{\Sigma}^{-1})_{ki} \quad (10a)$$

$$\hat{b}_i = \sum_k \hat{f}_k (R\tilde{\Sigma}^{-1})_{ki}, \quad (10b)$$

By construction these new particle operators also satisfy the Fermion anti-commutation relations, $\{\hat{b}_i^\dagger, \hat{b}_j\} = \delta_{ij}$. By inverting relations (10a)-(10b), using the expression (5a)-(5b) relating the $(\hat{\chi}_k^\dagger, \hat{\chi}_k)$ to the $(\hat{f}_k^\dagger, \hat{f}_k)$ and using (8), these new particle operators can be related to the q.p. operators $(\hat{\gamma}_\mu^\dagger, \hat{\gamma}_\nu)$ through

$$\hat{\gamma}_\mu^\dagger = \sum_l \hat{b}_l^\dagger \left[\tilde{\Sigma} R^\dagger \tilde{u} Q \Sigma^{-1} \right]_{l\mu} - \hat{b}_l \left[\tilde{\Sigma} R^\dagger \tilde{v} Q \Sigma^{-1} \right]_{l\mu} \quad (11a)$$

$$\hat{\gamma}_\mu = \sum_l \hat{b}_l^\dagger \left[\tilde{\Sigma} R^\top \tilde{u} Q^* \Sigma^{-1} \right]_{l\mu} + \hat{b}_l \left[\tilde{\Sigma} R^\top \tilde{v} Q^* \Sigma^{-1} \right]_{l\mu}. \quad (11b)$$

39 These two equations are the main result of this appendix. They show that we can extract from the non-
40 orthogonal reconstructed, canonical wavefunctions a set of quasiparticle operators that obey the Fermion
41 anticommutation relation, define a quasiparticle vacuum and are related to an orthonormal basis of the
42 single-particle Hilbert space through the following Bogoliubov transformation

$$\tilde{W} = \begin{pmatrix} \tilde{\Sigma} R^\top \tilde{u} Q^* \Sigma^{-1} & \tilde{\Sigma} R^\dagger \tilde{v} Q \Sigma^{-1} \\ \tilde{\Sigma} R^\top \tilde{v} Q^* \Sigma^{-1} & \tilde{\Sigma} R^\dagger \tilde{u} Q \Sigma^{-1} \end{pmatrix} = \begin{pmatrix} \tilde{U} & \tilde{V}^* \\ \tilde{V} & \tilde{U}^* \end{pmatrix}. \quad (12)$$

43 This matrix only depends on the initial canonical occupations \tilde{u} and \tilde{v} , as well as on the eigenvalues and
44 eigenvectors of both the s.p. overlap matrix (3) and the q.p. overlap matrix (6).

45 From the new transformation \tilde{W} , we can define the one-body density matrix

$$\rho = \tilde{V}^* \tilde{V}^\top = \tilde{\Sigma} R^\dagger \tilde{v} \tilde{\tau}^{(xx)^{-1}} \tilde{v} R \tilde{\Sigma}. \quad (13)$$

46 and put it into canonical form by diagonalizing it

$$\rho = \tilde{D} v^2 \tilde{D}^\dagger. \quad (14)$$

47 The transformation \tilde{D} defines the new canonical basis. By construction, these new canonical wavefunctions
 48 are expressed in the $(\hat{b}, \hat{b}^\dagger)$ basis, which is itself related to the original, non-orthogonal basis of the
 49 reconstructed wavefunctions $(\hat{f}, \hat{f}^\dagger)$ through (10a)-(10b). One can easily show that we have

$$\tilde{D}_n(\mathbf{r}, \sigma) = \sum_k \phi_k(\mathbf{r}, \sigma) (R\tilde{\Sigma}^{-1}\tilde{D})_{kn}. \quad (15)$$

50 At this point, we have obtained a set of genuine canonical wavefunctions $\tilde{D}_n(\mathbf{r}, \sigma)$ that are orthonormal
 51 and are associated with the new occupations v_n defined by (14). The relation between these canonical
 52 wavefunctions and the wavefunctions reconstructed by the AE is given by (15). Thanks to this expression,
 53 we can now apply the Haider & Gogny formulas for the norm overlap between two many-body states $|\Phi\rangle$
 54 and $|\Psi\rangle$. We find

$$\langle \Phi | \Psi \rangle = \det(\Sigma^{(\Phi)})^{-1} \det(\Sigma^{(\Psi)})^{-1} \det(\tau^{\Phi\Psi}) \det(\tilde{Z}^{(\Phi\Psi)}), \quad (16)$$

where

$$\tau_{mn}^{(\Phi\Psi)} = \sum_\sigma \int d^3\mathbf{r} \tilde{D}_m^{(\Phi)*}(\mathbf{r}, \sigma) \tilde{D}_n^{(\Psi)}(\mathbf{r}, \sigma), \quad (17a)$$

$$Z^{(\Phi\Psi)} = u^{(\Phi)} (\tau^{(\Phi\Psi)\dagger})^{-1} u^{(\Psi)} + v^{(\Phi)} \tau^{(\Phi\Psi)} v^{(\Psi)}, \quad (17b)$$

$$u^{(\Phi/\Psi)} = \sqrt{1 - v^{(\Phi/\Psi)2}}. \quad (17c)$$

2 METRIC INDUCED BY AN INNER PRODUCT

55 We present in this section the different notions of distance associated with an inner product. We note $\langle \mathbf{a} | \mathbf{b} \rangle$
 56 the inner product between two vectors \mathbf{a} and \mathbf{b} . There are many examples of inner product in nuclear
 57 physics, such as the overlap $\langle \Phi | \Psi \rangle$ between two many-body states $|\Phi\rangle$ and $|\Psi\rangle$ or the overlap between
 58 single-particle orbitals $\varphi(\mathbf{r}, \sigma)$ and $\phi(\mathbf{r}, \sigma)$ defined as

$$\langle \varphi | \phi \rangle = \sum_\sigma \int d^3\mathbf{r} \varphi^*(\mathbf{r}, \sigma) \phi(\mathbf{r}, \sigma). \quad (18)$$

59 Let us first recall some standard mathematics notations. The norm induced by the inner product $\langle \mathbf{a} | \mathbf{b} \rangle$ is
 60 defined in the usual way as

$$\|\mathbf{a}\| = \sqrt{\langle \mathbf{a} | \mathbf{a} \rangle}. \quad (19)$$

61 We can then introduce the distance induced by the inner product as

$$d_I(\mathbf{a}, \mathbf{b}) = \|\mathbf{a} - \mathbf{b}\|. \quad (20)$$

62 Note that this distance depends on the possible phase and norm of \mathbf{a} and \mathbf{b} . However, quantum-mechanical
 63 observables do not depend on either of them. The norm-independent distance thus reads

$$d_o(\mathbf{a}, \mathbf{b}) = \left\| \frac{\mathbf{a}}{\|\mathbf{a}\|} - \frac{\mathbf{b}}{\|\mathbf{b}\|} \right\|, \quad (21)$$

64 which can be rewritten as a function of the inner product between the two normed vectors

$$d_o(\mathbf{a}, \mathbf{b}) = \sqrt{2} \sqrt{1 - \Re \left[\left\langle \frac{\mathbf{a}}{\|\mathbf{a}\|} \middle| \frac{\mathbf{b}}{\|\mathbf{b}\|} \right\rangle \right]} = \sqrt{2} \sqrt{1 - \left| \left\langle \frac{\mathbf{a}}{\|\mathbf{a}\|} \middle| \frac{\mathbf{b}}{\|\mathbf{b}\|} \right\rangle \right| \cos \Theta}, \quad (22)$$

65 where $\Theta = \arg \langle \mathbf{a} | \mathbf{b} \rangle$. Another choice for a norm-invariant distance is the Great-Circle distance, also
 66 known as orthodromic or spherical distance, that is defined as Deza and Deza (2009)

$$d_{\perp}(\mathbf{a}, \mathbf{b}) = \arccos \left(\Re \left[\left\langle \frac{\mathbf{a}}{\|\mathbf{a}\|} \middle| \frac{\mathbf{b}}{\|\mathbf{b}\|} \right\rangle \right] \right) = \arccos \left(\left| \left\langle \frac{\mathbf{a}}{\|\mathbf{a}\|} \middle| \frac{\mathbf{b}}{\|\mathbf{b}\|} \right\rangle \right| \cos \Theta \right) \quad (23)$$

The orthodromic distance is defined on the manifold of unit vectors. In the case of real vector spaces, it can be interpreted as the angle between \mathbf{a} and \mathbf{b} . Equations (22) and (23) clearly show that both distances d_o and d_{\perp} still depend on the phase Θ between \mathbf{a} and \mathbf{b} . To remove this dependency, we minimize each distance d_o and d_{\perp} over Θ . This gives the following two norm- and phase-independent distances ²,

$$D(\mathbf{a}, \mathbf{b}) = \sqrt{2} \sqrt{1 - \left| \left\langle \frac{\mathbf{a}}{\|\mathbf{a}\|} \middle| \frac{\mathbf{b}}{\|\mathbf{b}\|} \right\rangle \right|}, \quad (24a)$$

and

$$D_{\perp}(\mathbf{a}, \mathbf{b}) = \arccos \left| \left\langle \frac{\mathbf{a}}{\|\mathbf{a}\|} \middle| \frac{\mathbf{b}}{\|\mathbf{b}\|} \right\rangle \right|. \quad (25a)$$

67 The distance D_{\perp} is an intrinsic metric and is named the Fubini–Study metric. It is a generalization of the
 68 Bloch sphere. Table 1 presents all the distances on the 1-body Hilbert space between orbitals that we have
 69 considered in this work.

Notation	Space	Invariance		
		Norm	Phase	Definition
d_I	\mathcal{H}	No	No	(20)
d_o	Unit vectors of \mathcal{H}	Yes	No	(21)
d_{\perp}	Unit vectors of \mathcal{H}	Yes	No	(23)
\bar{D}	Riemann sphere of \mathcal{H}	Yes	Yes	(24)
D_{\perp}	Riemann sphere of \mathcal{H}	Yes	Yes	(25)

Table 1. Different metrics can be defined on the set of orbitals.

70 All these distances are defined on the one-body Hilbert space of s.p. wavefunctions. As a result, they do
 71 not depend on the occupation probability of canonical orbitals, in contrast to the many-body state which
 72 takes the BCS form. As already mentioned in the main text, determining such a dependency exactly from
 73 (32) in Sec. 4.1.3 is not trivial and computationally demanding. Instead, we can adopt the approximation
 74 that the dependency should be proportional to some power p of the occupation number v_{μ}^2 associated with
 75 the current orbital,

$$d^{(p)}(\varphi, \phi) = (v_{\mu}^2)^p \times d(\varphi, \phi). \quad (26)$$

² Note that they are distances over the projective space $P(\mathcal{H})$, not over the 1-body Hilbert space \mathcal{H}

REFERENCES

- 76 Haider Q, Gogny D. Microscopic approach to the generator coordinate method with pairing correlations and
77 density-dependent forces. *J. Phys. G: Nucl. Part. Phys.* **18** (1992) 993. doi:10.1088/0954-3899/18/6/003.
- 78 Ring P, Schuck P. *The Nuclear Many-Body Problem*. Texts and Monographs in Physics (Springer) (2004).
- 79 Schunck N. *Energy Density Functional Methods for Atomic Nuclei*. IOP Expanding Physics (Bristol, UK:
80 IOP Publishing) (2019).
- 81 Deza M, Deza E. *Encyclopedia of Distances* (Springer) (2009). doi:10.1007/978-3-642-00234-2.