

# Analiza pjesme grbavog kita s Istočnih Kariba

---

Nikšić, Sara

Master's thesis / Diplomski rad

2014

Degree Grantor / Ustanova koja je dodijelila akademski / stručni stupanj: **University of Zagreb, Faculty of Science / Sveučilište u Zagrebu, Prirodoslovno-matematički fakultet**

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:217:340079>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-03-09**



Repository / Repozitorij:

[Repository of the Faculty of Science - University of Zagreb](#)



Sveučilište u Zagrebu  
Prirodoslovno-matematički fakultet  
Biološki odsjek

Sara Nikšić

**Analysis of humpback whale song  
from the Eastern Caribbean**

Diplomski rad

Zagreb, 2014

Ovaj rad je izrađen u School of Biology, University of St Andrews, United Kingdom, pod vodstvom Dr. Luke Rendell-a. Rad je predan na ocjenu Biološkom odsjeku Prirodoslovno-matematičkog fakulteta Sveučilišta u Zagrebu radi stjecanja zvanja Magistra Ekologije i zaštite prirode.

## **Acknowledgements**

First of all, I want to thank Luke Rendell for giving me this project, and for all his invaluable help and guidance. I couldn't hope for a better supervisor!

I would also like to thank Ivana Maguire for all her support, even with my stubbornness about the title of this project.

Next I want to thank Vincent Janik, who opened for me the door of an amazing world of marine mammal science, and Thomas Gotz for our little chats about how to survive in it.

I would like to express my gratitude to Danielle Cholewiak and Michael Noad for sharing their songs with me, and to Ellen Garland, Anita Murray, Elisa Girola and Jennifer Allen, who were already familiar with all the peculiarities of singing whales and took time to help me with their advice.

There are no words to describe how grateful I am to Simon Moss, one of the nicest and craziest people I have ever met, not just for letting me into his house, but also for making me feel at home.

Big thanks to Chris for always being there for me, especially all the times I couldn't make it to Simon's house.

To all the rest of my St Andrews friends, especially guys from SMRU and the Onsley family, I want to say thanks so much for making my time in The Bubble absolutely amazing! I will miss you all!

To the guys back home, I could write a book about all the good times we had during our studies, and even a bigger one about the fun we had beside the studying part. For now, I will just say thank you all for being my friends and making these years the best of my life!

Cimi, although there were so many times in these 4 years we lived together that I wanted to kick you out, I am glad I didn't. You also had to deal with me, which was not always easy, so thank you for that!

Hrvoje, I can't even put into words what I want to say to you, but I think you already know it. Thank you for being there for me even when I went chasing seals and whales!

Last but not least, I want to thank my family that supported all of my crazy dreams – I couldn't do it without you! You helped me become a person I am today, and whether that is a good or a bad thing, you still love me and thank you so much for that!

# TEMELJNA DOKUMENTACIJSKA KARTICA

Sveučilište u Zagrebu

Prirodoslovno-matematički fakultet

Biološki odsjek

Diplomski rad

## ANALIZA PJESME GRBAVOG KITA S ISTOČNIH KARIBA

Sara Nikšić

Rooseveltov trg 6, 10000 Zagreb, Hrvatska

Mužjaci grbavih kitova proizvode hijerarhijski strukturirane i kompleksne pjesme. Ove pjesme su naučene i konstantno se mijenjaju kroz evolucijske i revolucijske procese vodoravnog prijenosa kulture. Populacije iz istog oceana pokazuju sličnosti u svojim pjesmama, dok geografski odvojene populacije imaju potpuno različite pjesme. Iako su detaljne analize varijacija unutar pjesama istih i različitih jedinki rijetke, prijeko su potrebne za razumijevanje kako individualno učenje pjesama utječe na kulturne promjene na razini populacija, koje se proučavaju od 1970-ih godina. U ovom istraživanju analizirane su pjesme grbavih kitova iz sjevernog Atlantika i južnog Pacifika. Za klasifikaciju jedinica pjesme je testirana automatizirana metoda, ali, budući da se pokazala nezadovoljavajućom, podjela pjesmi se oslanjala na ljudsku procjenu. Pjesme su podijeljene na jedinice, fraze i teme, koje su onda uspoređene metodom Levenshteinove udaljenosti. Rezultati su pokazali da pjesme iz sjevernog Atlantika dijele značajan dio materijala, ali da pjesma iz Australije ne pokazuje nikakve sličnosti na razini fraza i tema s pjesmama iz sjevernog Atlantika. Ovaj rad pruža dokaze o vodoravnom prijenosu pjesama grbavih kitova unutar istog oceana, koji se ne događa između geografski udaljenih područja.

(34 stranica, 11 slika, 9 tablica, 49 literaturnih navoda, jezik izvornika: engleski)

Rad je pohranjen u Središnjoj biološkoj knjižnici

Ključne riječi: kitova pjesma, prijenos kulture, vokalna komunikacija, Levenshteinova udaljenost

Voditelj: Dr. Luke Rendell

Suvoditelj: Izv. prof. dr. sc. Ivana Maguire

Ocjenitelji: Izv. prof. dr. sc. Ivana Maguire, Doc. dr. sc. Duje Lisičić, Izv. prof. dr. sc. Antun Alegro, Doc. dr. sc. Jasna Lajtner

Rad prihvaćen: 8. 5. 2014.

## BASIC DOCUMENTATION CARD

University of Zagreb

Faculty of Science

Division of Biology

Graduation Thesis

### **ANALYSIS OF HUMPBACK WHALE SONG FROM THE EASTERN CARIBBEAN**

Sara Nikšić

Rooseveltova trg 6, 10000 Zagreb, Croatia

Male humpback whales produce hierarchically structured and complex songs. These songs are learned, and are constantly changing at both evolutionary and revolutionary tempos through horizontal cultural transmission processes. Populations within the same ocean basin share similarities in their song displays, whereas geographically distinct populations have completely different songs. However, fine-scale analyses of variation in song production within and between individuals are sparse, but essential for understanding the details of how individual song learning can produce the population level cultural changes that have been observed since the 1970s. In this study, songs of humpback whales from the North Atlantic and South Pacific Ocean were analysed. An automated method for song unit classification was tested, against classification by a human observer, and the latter proved more effective. All songs were divided into unit, phrase and theme sequences which were then compared using the Levenshtein distance metric. The results showed that songs from North Atlantic shared a significant portion of their material, and that the Australian song had no similarity at phrase and theme level with any of the North Atlantic songs. This study provides evidence of horizontal cultural transmission of humpback whale songs within the same ocean basin that does not occur with a wider geographical separation.

(34 pages, 11 figures, 9 tables, 49 references, original in: English)

Thesis deposited in the Central Biological Library

Key words: whale song, cultural transmission, vocal communication, Levenshtein distance

Supervisor: Dr. Luke Rendell

Co-supervisor: Assoc. Prof. Ivana Maguire

Reviewers: Assoc. Prof. Ivana Maguire, Assist. Prof. Duje Lisičić, Assoc. Prof. Antun Alegro, Assist. Prof. Jasna Lajtner

Thesis accepted: 8. 5. 2014.

# Contents

Introduction.....	1
Aims.....	4
Hypothesis.....	4
Materials and methods .....	5
Recordings of songs .....	5
Acoustic analysis.....	6
Classification of units .....	6
Classification of phrases and themes.....	8
Song classification.....	9
The ‘splitting’ and ‘lumping’ issue .....	10
Transcription of names.....	12
Levenshtein distance analysis .....	14
Results.....	15
Principal component analysis (PCA) .....	15
The structure of songs .....	17
Levenshtein distance analysis .....	20
Discussion.....	25
Review of the methods used.....	25
Unit classification .....	25
Delineation of phrases and themes .....	25
Levenshtein distance method.....	26
Comparison with the previous studies .....	26
Future research .....	27
Conclusions.....	28
References.....	29
Appendices.....	34



## Introduction

There are many different ways in which animals communicate, but few can compete with humpback whales (*Megaptera novaeangliae*) when it comes to the complexity of their vocal displays. Humpback whales can produce a variety of sounds that range from ‘social sounds’ that both males and females generate (Winn *et al.* 1979, Tyack 1983, Dunlop *et al.* 2007), to complex and highly structured songs (Payne & McVay 1971, Winn & Winn 1978, Payne *et al.* 1983) which are sung just by males (Darling *et al.* 1983, Glockner 1983).

Humpbacks belong to the baleen whales, or mysticetes, which have baleen instead of teeth, and two blowholes, opposed to toothed whales which have just one. They can grow up to 16 m and weigh more than 40 tons, but the average length for females is 13 m, and for males 12.5 m. Both females and males reach sexual maturity between the ages of 5 and 10, although males don’t start breeding until later. Females give birth every two to three years to a single calf, which they nurse for a year. Humpback whales have a cosmopolitan distribution. Except the Arabian Sea population that remains there for the entire year, humpbacks from all around the world undertake long seasonal migrations from high latitude summer feeding grounds to low latitude winter feeding grounds. There are an estimated 35 – 40 000 humpbacks in the Northern Hemisphere and about 50 000 in the Southern hemisphere (Clapham 2013).

In 1971, Payne and McVay established the terminology for hierarchical structure of a humpback whale song, and the same classification of song elements has been used ever since. The smallest element of a song is a unit and it is defined as ‘the shortest sound that is continuous to our ears when heard in real time’ (Payne & McVay 1971). A repeated sequence of units comprises a phrase, several phrases make up a theme and a few themes constitute a song. Male humpbacks often cyclically repeat their songs during bouts of production, each repetition is referred to as a song cycle, and one or more song cycles make up a song session. These sessions can last for hours; Winn & Winn (1978) reported one whale singing continuously for 22 h. Although there is clearly some consistency in the song structure, there is quite a bit of variability both within- and between- individual whale songs.

Humpbacks from around the world have the same repertoire of units they can produce, but the order in which they are arranged varies in different areas (Payne and Guinee 1983). All individuals within a breeding population sing a similar version of the song, and populations

from the same ocean basin share similarities in their song displays, but geographically distinct populations have completely different songs (Winn *et al.* 1981, Payne & Guinee 1983, Payne *et al.* 1983, Helweg *et al.* 1998, Cerchio *et al.* 2001, Darling & Sousa-Lima 2005).

A number of studies have described changes in songs across populations (Winn *et al.* 1981, Guinee *et al.* 1983, Payne *et al.* 1983, Payne & Guinee 1983, Helweg *et al.* 1998, Cerchio *et al.* 2001), but variation within the same population has also been documented. Some songs of the individual whales were so different from the ‘norm’ of the particular temporal and spatial region that Frumhoff (1983) termed them ‘aberrant’.

Male humpbacks sing predominantly on the winter breeding grounds (Payne & McVay 1971, Winn & Winn 1978) and on the migration routes between breeding and summer feeding grounds (Payne & McVay 1971, Clapham & Mattila 1990), but they are also heard singing on their feeding grounds (Mattila *et al.* 1987, McSweeney *et al.* 1989, Clark & Clapham 2004).

Since it appears that only males sing, sexual selection is a leading hypothesis for the evolution of humpback song. However, while different hypotheses have been proposed about the role of singing in the humpback whale mating system, it is still unclear whether the primary function is of inter- or intra- sexual nature. Some researchers suggest that the songs have a role in female attraction (Tyack 1981, Clapham 1996, Smith *et al.* 2008), while others believe that its purpose is largely related to male – male interactions (Darling *et al.* 2006, Cholewiak 2008). Song may of course function in both ways.

Researchers noticed early on that songs slowly evolve from one form to another (Guinee *et al.* 1983, Payne *et al.* 1983, Payne & Payne 1985), but later research has also revealed that one populations’ song can be completely replaced by a song from the other population within a period of just two years, which was referred to as ‘cultural revolution’ (Noad *et al.* 2000). This was initially reported as occurring just once in the Eastern Australia breeding population (Noad *et al.* 2000), but it has since been shown to be a regular occurrence in populations all across the South Pacific, which appear to regularly discard their own song completely and instead take up the song of a neighbouring population *in toto* (Garland *et al.* 2011). Both entire songs and song changes are rapidly spread across the populations via vocal learning through a process of horizontal (i.e. largely within-generation) cultural transmission (Rendell & Whitehead 2001, Eriksen *et al.* 2005, Garland *et al.* 2011).

Payne and Guinne (1983) proposed three potential paths for exchanging songs between different populations. The first one was that it might happen inbetween breeding seasons on common migration routes and/or feeding grounds, for which Garland *et al.* recently (2013) provided evidence. The second possibility was that individuals might go to more than one breeding ground within the same season, and the third one was that individuals go to different breeding grounds in subsequent winters. However, these last two possibilities are unlikely, because we now know that humpbacks show strong maternally determined fidelity to breeding grounds (Clapham *et al.* 1993, Palsbøll *et al.* 1995, Stevick *et al.* 2006).

Analysing the songs of humpbacks involves one of the most fundamental issues in studying animal behaviour, the categorisation of behavioural patterns (Janik 1999). In bioacoustics, the problem of delineating appropriate categories for sounds is perennial, whether talking about sperm whale codas (Rendell & Whitehead 2003), signature whistles in dolphins or killer whale calls (Deecke & Janik 2006). Because the same problem exists with delineation of humpback whale song (Cholewiak *et al.* 2013), there have been several attempts to develop automated methods for classifying song elements in the last decade (Maeda *et al.* 2000, Suzuki *et al.* 2006, Green *et al.* 2010, Pace *et al.* 2010, Garland *et al.* 2012). Recently, Cholewiak *et al.* (2013) carried out an extensive review of current classification issues in which they acknowledge that ‘no system has yet been demonstrated which compares with the pattern recognition skills of the human brain’ and propose a set of guidelines for consistent delineation of songs.

I analysed three song recordings from different locations: two from the North Atlantic (Caribbean Island of Dominica and Georges Bank) and one from the Southwest Pacific (off of Peregian Beach, Queensland). The Antillean Island chain (which includes Dominica) is a known breeding area for humpbacks and Georges Bank is one of their North Atlantic feeding grounds (Kennedy *et al.* 2014). Humpback whales from Southwest Pacific migrate along the eastern coast of Australia, so they pass Peregian beach twice every year while migrating between their feeding and breeding grounds (Cato 1991, Noad & Cato 2001).

## **Aims**

The aims of this study were to:

1. Conduct a detailed analysis of the song recorded near the Caribbean Island of Dominica using both manual and automated method for unit classification
2. Decide which method for classifying units is to be used for the other songs
3. Delineate songs into unit, phrase and theme sequences
4. Estimate how many song units are shared within and between populations
5. Explain how different sequences are distributed across songs
6. Compare the structure of all songs at unit, phrase and theme level using the Levenshtein distance method
7. Characterise individual variation in song production (how similar are the songs from the same song session of the same whale)
8. Examine how much variance is there between different whales' song repertoires, both within the same ocean basin and between different oceans

## **Hypothesis**

The hypothesis was, according to the current literature on humpback song, that songs from the North Atlantic should share some similarities between themselves, but very few with the Australian song (with the possibility of sharing some parts of the unit repertoire).


## Materials and methods

### Recordings of songs

The first song was recorded on 9<sup>th</sup> of May 2012 at 14:11:48 using a towed 2-element hydrophone array based on Benthos AQ-4 elements (Schulz *et al.* 2011) deployed from a 14 m sailboat near the Caribbean Island of Dominica (15° N, -61° W) (Figure 1). The recording was made on a laptop PC using a Fireface 400 sound card and Pamguard software with a sampling rate of 96000 Hz.

The second song was recorded in Georges Bank (40° N, -69° W) (Figure 1) on 17<sup>th</sup> of April 2012 at 12:45:00 using Marine Autonomous Recording Units (Calupca *et al.* 2000) with sampling rate of 5000 Hz. The relatively low sampling rate for this recording meant that only frequencies up to 2500 Hz were registered.

The third song was recorded on 9<sup>th</sup> of September 2012 at 7:37:32 with an Acousonde 3A recorder (Sousa-Lima *et al.* 2013) deployed off of Peregian Beach, Queensland (-26° N, 153° E) (Figure 1). The sample rate was set to 27330 Hz, but an anti-aliasing filter was used that restricted the maximum frequency to 9292 Hz, so again the high frequencies were cut off.




**Figure 1:** Locations of the recording sites

## Acoustic analysis

Regardless of the quality of recordings (which varied considerably in both sampling rates and signal-to-noise ratios), all songs were viewed as spectrograms in Adobe Audition 3.0, calculated using a Blackmann–Harris window and Fast Fourier Transform (FFT) size of 1024. The songs were split into themes, phrases and units based on their visual and aural characteristics, taking into account the highly structured hierarchy of the song and the position of song elements in the song. The delineation of songs can be very challenging and subjective, so guidelines presented by Cholewiak *et al.* (2013) were followed.

## Classification of units


The first song was divided into units using markers in Adobe Audition 3.0 (Figure 2). Units were classified into types and units of the same type were copied into the same file. Every file was imported separately in Raven Pro 1.3 for further analysis.



**Figure 2:** Example of unit markers in Adobe Audition 3.0

Spectrograms of units were produced in Raven Pro 1.3 using a Hamming window and FFT size of 8192. Measurements were made using range selections (Figure 3). By default, Raven performs four measures that define the edges of selections in spectrogram view – begin time, end time, low frequency and high frequency. In addition to these default measures, maximum

frequency (the frequency at which maximum power occurs within the selection), delta frequency (the difference between the upper and lower frequency limits of the selection) and delta time (the difference between begin and end time for the selection) were also calculated, as well as robust signal measurements (measurements that do not vary much with changing the selection borders). The robust measurements included: center frequency (the frequency that divides the selection into two frequency intervals of equal energy), first quartile frequency (the frequency that divides the selection into two frequency intervals containing 25% and 75% of the energy in the selection), third quartile frequency (the frequency that divides the selection into two frequency intervals containing 75% and 25% of the energy in the selection) and inter-quartile range bandwidth (the difference between the first and the third quartile frequency).




**Figure 3:** Example of range selections in Raven Pro 1.3

In order to test if the measurements taken in Raven Pro 1.3 could explain the human classification of units, a principal component analysis (PCA) was conducted in Matlab R2013b. It turned out that the principal components were not enough to explain all the variation within the unit types, as will be shown in the results. The decision was made to retain the original classification and use only a human classifier for the rest of the songs.

The songs from the second and the third recording were viewed as spectrograms in Adobe Audition 3.0 and each unit was given a descriptive name (e. g., moan, groan, squeak, whoop) to make the process of unit recognition faster. Descriptive names were then replaced with coded names (e. g., a, b, c, d). There were seven unit types that occurred in more than one song, so the same names were used for the units of the same type, regardless of which song they came from. Spectrograms of all the unit types are presented in Appendix I and all song sequences represented by unit codes are shown in Appendix II.

### **Classification of phrases and themes**


The classification of phrases and themes was especially challenging because there were many ways in which they could be delineated. However, as many authors (Frumhoff 1983, Payne *et al.* 1983, Cerchio *et al.* 2001) agreed that phrase duration is one of the most stable characteristics of the humpback whale song, duration was the main criterion used for the phrase interpretation. Phrases were measured from the beginning of one to the beginning of another (subsequent) phrase, as suggested by Cholewiak *et al.* (2013). Figure 4 shows two different phrase types, both lasting for 14 seconds.



**Figure 4:** Duration of phrases



Because the number of phrases in themes can be extremely variable, the duration of themes is not a reliable feature, therefore decisions about theme delineation depended only on theme structure. Figure 5 shows the end of one and the beginning of the other theme.



**Figure 5:** The switch from one theme to another

### **Song classification**

The Eastern Caribbean recording captured only one complete song cycle, so the analysis included just one song from that area which was named ‘song 1’. The recording from Georges Bank included six full songs from the same song session and all of them were analysed. They were given names that started with the number two followed by consecutive letters (e.g., 2a, 2b, 2c). The Australian recording contained part of a song session which included three songs. Because the first one was not recorded from the beginning and the recording stopped before the end of the third one, just the second song (named ‘song 3’) was included in the analysis.

## **The ‘splitting’ and ‘lumping’ issue**

The biggest challenge in delineating songs was to decide how similar two elements of a song have to be to be treated as of the same type, or how different should they be to be assigned different names. In the extreme case, ‘splitter’ approach would be the one in which a new type is assigned every time there is even a small difference, and ‘lumper’ would be if everything that is remotely similar was grouped together.

In songs 1 and 2 some units evolved throughout the song and changed slowly from one type to another, so the problem was to decide at which point in the song the unit changed enough to allocate a new name. The Australian recording was much easier to deal with regarding the unit classification, because the units did not alter during the entire song. For the songs 1 and 2 I tried to decide on something in-between the two approaches, so I allowed some degree of variation in frequency and duration of units, but was very rigorous about the position of units within phrases.

The ‘splitting’ and ‘lumping’ issue was also present when classifying phrases and themes. They also changed throughout the songs; for example, the same unit could be repeated a few more times in subsequent phrases of the same type, or the number of phrase repetitions within themes of the same type could also vary; sometimes one unit stopped appearing, or a new one was introduced, without changing the overall pattern of the song. These examples are shown from a ‘lumper’ perspective; a ‘splitter’ would name all of them different types. Example of theme classification using two different approaches is shown in Figure 6.

SPLITTING	
song	themes
1	1Aa-1Ab-1B-1Ca-1Cb-1Da-1Db-1Ea-2A-2B-2C-2D-3-4
2a	5Aa-5Ba-5Bb-5Bc-1Fa-1Cc-1Ga-1Ha
2b	5Ab-5Ac-5Ad-5Ae-5Af-5Ac-5Bb-5Bd-1Fb-1Fc-1Fd-1Fd-1Fe-1Ff-1Cd-1Gb-1Hb
2c	5Ac-5Ac-5Af-5Be-5Bf-5Bg-1Fg-1Fd-1Fb-1Fh-1Cd-1Ce-1Ce-1Gc-1Hc
2d	5Ac-5Ac-5Ag-5Ac-5Bh-5Be-5Bj-5Bc-1Fi-1Fc-1Fj-1Ff-1Fk-1Fj-1Cf-1Ce-1Dc-1Dd-1De-1Eb-1Hd
2e	5Af-5Ah-5Ac-5Ca-5Cb-5Cc-5Cd-5Ce-5Cf-5Bc-5Bk-1Fb-1Fl-1Ff-1Fj-1Fm-1Cg-1Ch-1Gd-1He
2f	5Cg-5Ch-5Ci-5Ai-5Ab-5Bl-5Bm-5Be-5Bn-5Bc-1Fi-1Ff-1Fn-1Fm-1Fm-1Ci-1Ce-1Ec-1Hf
3	6-7-8-9-10-11
LUMPING	
song	themes
1	1-1-1-1-1-1-1-1-2-2-2-2-3-4
2a	5-5-5-5-1-1-1-1
2b	5-5-5-5-5-5-5-5-1-1-1-1-1-1-1-1
2c	5-5-5-5-5-5-1-1-1-1-1-1-1-1
2d	5-5-5-5-5-5-5-5-1-1-1-1-1-1-1-1-1-1-1
2e	5-5-5-5-5-5-5-5-5-5-1-1-1-1-1-1-1-1
2f	5-5-5-5-5-5-5-5-5-5-1-1-1-1-1-1-1-1
3	6-7-8-9-10-11

**Figure 6:** ‘Splitting’ and ‘lumping’ of theme strings

## **Transcription of names**

The sequences (or strings) of phrases were named with simple combinations of letters and numbers (e.g., A1a, A1b, B1a, B1b). Similar transcription was carried out for themes, but the combinations of characters used were different (e.g., 1Aa, 1Ab) in order to make it more transparent.

For example, the song 1 can be represented by a theme string:

1Aa-1Ab-1B-1Ca-1Cb-1Da-1Db-1Ea-2A-2B-2C-2D-3-4;

and theme 1Aa can be represented by sequence of phrases:

A1a-A1a-A1a-A1a-A1a-A1b-A1c-A1c-A2-B.

As with the units, the same phrases and themes (regardless of which song they came from) were given the same names.

They were then converted into coded names which were later used in the Levenshtein distance analysis. Coding was done so that the same letters and numbers could be used just within the same phrase type, because the results of the analysis were incorrect if just different combinations of the same characters were used. Table 1 contains all the original and corresponding coded names of themes and phrases. Similar colours were used for similar themes and phrases (e. g., blue for themes of type 1) and black was used for all the themes and phrases that came from the Australian song because none of them occurred in any other songs.

**Table 1:** Transcription of names

original names		coded names		original names		coded names	
themes	phrases	themes	phrases	themes	phrases	themes	phrases
IAa	A1a	IAa	A1a	2A	Ib	2Π	IIw
IAb	A1b	IAb	A1b	2B	Ic	2JJ	IIç
IB	A1c	IBs	A1c	2C	Ja	2KK	JJr
ICa	A1d	ICc	A1d	2D	Jb	2LL	JJḡ
ICb	A2	ICd	A22	3	Jc	333	JJḡ
ICc	A3a	ICe	A3e	4	Ka	444	KKΠḡ
ICd	A3b	ICf	A3f	5Aa	Kb	5Mτ	KKø
ICe	A3c	ICg	A3g	5Ab	La	5Mυ	LLç
ICf	A3d	ICḡ	A3h	5Ac	Lb	5Mφ	LLγ
ICg	A3e	ICi	A3i	5Ad	Lc	5Mχ	LLḡ
ICḡ	A3f	ICj	A3j	5Ae	UNP	5Mψ	UNP
ICi	A3g	ICk	A3k	5Af		5Mω	
IDa	A4a	IDl	A4l	5Ag		5Mḡ	
IDb	A4b	IDm	A4m	5Ah		5Mβ	
IDc	A5	IDn	A55	5Ai		5Mγ	
IDd	A6a	IDo	A6n	5Ba		5Nδ	
IDe	A6b	IDp	A6o	5Bb		5Nḡ	
IEa	A6c	IEq	A6p	5Bc		5Nз	
IEb	A6d	IEr	A6q	5Bd		5Ni	
IEc	A6e	IEs	A6r	5Be		5Ni	
IFa	A6f	IFt	A6s	5Bf		5Nл	
IFb	B	IFu	BBB	5Bg		5Nm	
IFc	C1	IFv	CCt	5Bh		5Nн	
IFd	C2	IFw	CCu	5Bi		5Nπ	
IFe	C3	IFx	CCv	5Bj		5Nц	
IFf	C4	IFy	CCw	5Bk		5Nч	
IFg	C5	IFz	CCx	5Bl		5Nш	
IFḡ	C6	IFa	CCy	5Bm		5Nщ	
IFi	C7	IFβ	CCz	5Bn		5Nъ	
IFj	D	IFγ	DDD	5Ca		5Oь	
IFk	Ea	IFδ	EEk	5Cb		5Oэ	
IFl	Eb	IFe	EEḡ	5Cc		5Oю	
IFm	F1a	IFç	FGḡ	5Cd		5Oя	
IFn	F1b	IFη	FGḡ	5Ce		5Oђ	
IGa	F2a	IGθ	FIçw	5Cf		5Oе	
IGb	F2b	IGi	FIçḡ	5Cg		5Oљ	
IGc	F3a	IGκ	FΔḡ	5Ch		5Oњ	
IGd	F3b	IGλ	FΔḡ	5Ci		5Oћ	
IHa	F3c	IHμ	FΔḡ	6		666	
IHb	G	IHξ	GGG	7		777	
IHc	Ha	IHπ	HHḡ	8		888	
IHd	Hb	IHρ	HHø	9		999	
IHe	Hc	IHç	HHo	10		£££	
IHf	Ia	IHσ	IIç	11		\$\$\$	

## **Levenshtein distance analysis**

All the coded unit, phrase and theme strings were imported into Matlab R2013b, and using custom written code the unweighted Levenshtein distance analysis (LD) was undertaken. LD is a method that calculates the minimum number of insertions (i), deletions (d) and substitutions (s) needed for one string (str1) to convert into another (str2):

$$\mathbf{LD (str1, str2) = \min (i+d+s)}$$

The method used was the unweighted LD that doesn't differentiate between insertions, deletions and substitutions; all of them have the same value of 1, thus assuming that all types of change hold equal significance. Because the longer sequences produced bigger LD scores, the distance was normalised by the length of the longest string:

$$\mathbf{LD_{norm} (str1, str2) = \min (i+d+s) / \max [\text{length (str1), (str2)}]}$$

Every theme string was compared to all of the other theme strings from all songs. The same analysis was conducted for all the phrase and unit strings. We used average – linkage cluster analysis which produced the agglomerative hierarchical cluster trees that showed how closely related are the songs at the unit, phrase and theme level.

Because there is no agreement about which approach for delineating songs is better ('splitter' or 'lumper'), Levenshtein distance analysis was carried out for both.

# Results

## Principal component analysis (PCA)


The four unit measurements entered into the PCA were: center frequency, delta time, first quartile frequency and third quartile frequency. The first two principal components explained 87% of the variability in the data. In Table 2 columns represent principal components, rows represent unit variables and the cell values are the loading coefficients. The first principal component (PC1) is mostly loaded on frequency measurements, and the second principal component (PC2) is almost completely loaded on duration.

The results shown in Figure 7 suggested that the features other than these measurements have to be taken into account when classifying units. Different colours represent different unit types. Even though some of the units (e. g., unit ‘a’) were clearly separated from the others, most of the unit types were overlapping with the others (e. g., unit ‘d’ and unit ‘f’). The automated PCA lacked the ability to recognize one very important aspect of the song structure, and that is the position of units in the song. Because of that, the original classification by a human observer was maintained.

**Table 2:** Principal component loading coefficients

PC1, PC2, PC3, PC4: principal components

	<b>PC1</b>	<b>PC2</b>	<b>PC3</b>	<b>PC4</b>
<b>center frequency</b>	0.6101	-0.0372	-0.0417	0.7904
<b>delta time</b>	0.0303	0.9834	0.176	0.0322
<b>first quartile frequency</b>	0.5561	-0.1325	0.7176	-0.3977
<b>third quartile frequency</b>	0.5636	0.1182	-0.6725	-0.4649



**Figure 7:** The results of the principal component analysis

PC1: 1<sup>st</sup> principal component, PC2: 2<sup>nd</sup> principal component

a, b, c, d, e, f, g, h, i: unit types



## The structure of songs

All songs were divided into themes, phrases and units. Table 3 shows how many different types of themes, phrases and units there were in each song. Songs from Georges Bank had the smallest variation at all levels, so they were clearly the least complex of all songs.

**Table 3:** Number of theme, phrase and unit types

song	location	No. theme types	No. phrase types	No. unit types
1	Dominica	4	6	9
2a	Georges Bank	2	4	7
2b	Georges Bank	2	4	7
2c	Georges Bank	2	4	7
2d	Georges Bank	2	4	7
2e	Georges Bank	2	4	7
2f	Georges Bank	2	4	7
3	Peregrian Beach	6	6	12

Songs from Georges Bank shared a substantial proportion of their elements with the song from Dominica. One out of two themes present in the songs from Georges Bank also appeared in the Dominica song. The other theme was specific to Georges Bank (Table 4). The same pattern could be observed with phrases and units. Three out of four phrase types and six out of seven unit types from Georges Bank were present in Dominica song (Table 5 and Table 6). Other types of themes, phrases and units that were not seen in songs from Georges Bank were unique to Dominica song, except one unit type that was also present in the Australian song.

The Australian song shared no similarities with the songs from North Atlantic at the level of themes and phrases (Table 4 and Table 5). However, there was some similarity at the unit level. Unit type ‘g’ showed up in both Australian and Dominica song, and unit type ‘f’ was present in all songs (Table 6). The examples of unit ‘f’ from all three locations are represented in Figure 8.

**Table 4:** Theme types


theme	song 1	song 2a	song 2b	song 2c	song 2d	song 2e	song 2f	song 3
1	yes	yes	yes	yes	yes	yes	yes	no
2	yes	no	no	no	no	no	no	no
3	yes	no	no	no	no	no	no	no
4	yes	no	no	no	no	no	no	no
5	no	yes	yes	yes	yes	yes	yes	no
6	no	no	no	no	no	no	no	yes
7	no	no	no	no	no	no	no	yes
8	no	no	no	no	no	no	no	yes
9	no	no	no	no	no	no	no	yes
10	no	no	no	no	no	no	no	yes
11	no	no	no	no	no	no	no	yes

**Table 5:** Phrase types

phrase	song 1	song 2a	song 2b	song 2c	song 2d	song 2e	song 2f	song 3
A	yes	yes	yes	yes	yes	yes	yes	no
B	yes	yes	yes	yes	yes	yes	yes	no
C	yes	no	no	no	no	no	no	no
D	yes	yes	yes	yes	yes	yes	yes	no
E	yes	no	no	no	no	no	no	no
F	no	yes	yes	yes	yes	yes	yes	no
G	no	no	no	no	no	no	no	yes
H	no	no	no	no	no	no	no	yes
I	no	no	no	no	no	no	no	yes
J	no	no	no	no	no	no	no	yes
K	no	no	no	no	no	no	no	yes
L	no	no	no	no	no	no	no	yes

**Table 6:** Unit types

unit	song 1	song 2a	song 2b	song 2c	song 2d	song 2e	song 2f	song 3
<b>a</b>	yes	yes	yes	yes	yes	yes	yes	no
<b>b</b>	yes	yes	yes	yes	yes	yes	yes	no
<b>c</b>	yes	yes	yes	yes	yes	yes	yes	no
<b>d</b>	yes	no	no	no	no	no	no	no
<b>e</b>	yes	yes	yes	yes	yes	yes	yes	no
<b>f</b>	yes	yes	yes	yes	yes	yes	yes	yes
<b>g</b>	yes	no	no	no	no	no	no	yes
<b>h</b>	yes	no	no	no	no	no	no	no
<b>i</b>	yes	yes	yes	yes	yes	yes	yes	no
<b>j</b>	no	yes	yes	yes	yes	yes	yes	no
<b>k</b>	no	no	no	no	no	no	no	yes
<b>l</b>	no	no	no	no	no	no	no	yes
<b>m</b>	no	no	no	no	no	no	no	yes
<b>n</b>	no	no	no	no	no	no	no	yes
<b>o</b>	no	no	no	no	no	no	no	yes
<b>p</b>	no	no	no	no	no	no	no	yes
<b>q</b>	no	no	no	no	no	no	no	yes
<b>r</b>	no	no	no	no	no	no	no	yes
<b>s</b>	no	no	no	no	no	no	no	yes
<b>t</b>	no	no	no	no	no	no	no	yes



**Figure 8:** Unit 'f' from the song 1, song 2 and song 3

## Levenshtein distance analysis


Presented below are the results of the Levenshtein distance analysis. The value of 0 stands for the same strings; whereas the value of 1 means that the strings are completely different.

Table 7 and Figure 9 present the results of the LD analysis for the unit strings. LD values are shown in Table 7. Songs ‘2’ were the most similar within themselves, which was expected because they all came from the same song session. Song ‘1’ was the closest one to the songs ‘2’, and the song ‘3’ was the most distant one of all. It had the highest LD values compared to songs 1 and 2, but they were still not 1 because the Australian song did share some units with the other songs.

**Table 7:** Distances between unit strings

**UNITS**

	Song 1	Song 2a	Song 2b	Song 2c	Song 2d	Song 2e	Song 2f	Song 3
Song 1	0	0.774752	0.673267	0.690594	0.665842	0.678218	0.648515	0.876238
Song 2a	0.774752	0	0.462617	0.418367	0.528	0.5	0.531746	0.935484
Song 2b	0.673267	0.462617	0	0.214953	0.264	0.323009	0.313492	0.945161
Song 2c	0.690594	0.418367	0.214953	0	0.28	0.389381	0.357143	0.932258
Song 2d	0.665842	0.528	0.264	0.28	0	0.4	0.384921	0.945161
Song 2e	0.678218	0.5	0.323009	0.389381	0.4	0	0.261905	0.987097
Song 2f	0.648515	0.531746	0.313492	0.357143	0.384921	0.261905	0	0.993548
Song 3	0.876238	0.935484	0.945161	0.932258	0.945161	0.987097	0.993548	0



**Figure 9:** Distances between unit strings

The results of the LD analysis for phrase strings are shown in Table 7 and Figure 9. Because of the ‘splitting’ and ‘lumping’ issue, both approaches to phrase delineation were used. The LD values from the ‘lumper’ analysis were smaller than the ‘splitter’ values, but the overall pattern was the same. There was a substantial variation within the songs from Georges Bank, even though they came from the same song session of the same individual. The songs from Georges Bank shared some similarities with the Dominica song. All of the Australian song LD scores equalled 1, which means it was completely different from any other song.



**Table 7:** Distances between phrase strings

**PHRASES**

<b>SPLITTER</b>	Song 1	Song 2a	Song 2b	Song 2c	Song 2d	Song 2e	Song 2f	Song 3
Song 1	0	0.85977	0.806897	0.786207	0.783908	0.848276	0.848276	1
Song 2a	0.85977	0	0.537954	0.501792	0.556604	0.545455	0.564815	1
Song 2b	0.806897	0.537954	0	0.264026	0.292453	0.339934	0.32716	1
Song 2c	0.786207	0.501792	0.264026	0	0.320755	0.393939	0.361111	1
Song 2d	0.783908	0.556604	0.292453	0.320755	0	0.380503	0.373457	1
Song 2e	0.848276	0.545455	0.339934	0.393939	0.380503	0	0.287037	1
Song 2f	0.848276	0.564815	0.32716	0.361111	0.373457	0.287037	0	1
Song 3	1	1	1	1	1	1	1	0

<b>LUMPER</b>	Song 1	Song 2a	Song 2b	Song 2c	Song 2d	Song 2e	Song 2f	Song 3
Song 1	0	0.767123	0.650685	0.616438	0.671233	0.726027	0.732877	1
Song 2a	0.767123	0	0.49505	0.451613	0.518868	0.484848	0.527778	1
Song 2b	0.650685	0.49505	0	0.108911	0.103774	0.217822	0.157407	1
Song 2c	0.616438	0.451613	0.108911	0	0.141509	0.222222	0.222222	1
Song 2d	0.671233	0.518868	0.103774	0.141509	0	0.207547	0.185185	1
Song 2e	0.726027	0.484848	0.217822	0.222222	0.207547	0	0.12963	1
Song 2f	0.732877	0.527778	0.157407	0.222222	0.185185	0.12963	0	1
Song 3	1	1	1	1	1	1	1	0



**Figure 9:** Distances between phrase strings



The LD values for theme strings are shown in Table 8 and Figure 10. As with the phrases, both ‘splitter’ and ‘lumper’ analysis was conducted for the theme strings. Again, the LD values from the ‘splitter’ analysis were higher than the ones from the ‘lumper’ analysis. The songs from Georges Bank varied even more within themselves at the theme than at the phrase level. The LD scores of the song ‘1’ in comparison to songs ‘2’ were also higher than in the analysis of phrase strings. All of the LD scores of the Australian song were still 1.

**Table 8:** Distances between theme strings

**THEMES**

<b>SPLITTER</b>	Song 1	Song 2a	Song 2b	Song 2c	Song 2d	Song 2e	Song 2f	Song 3
Song 1	0	0.904762	0.941176	0.933333	0.888889	0.95	0.947368	1
Song 2a	0.904762	0	0.686275	0.644444	0.746032	0.75	0.719298	1
Song 2b	0.941176	0.686275	0	0.431373	0.47619	0.483333	0.508772	1
Song 2c	0.933333	0.644444	0.431373	0	0.492063	0.533333	0.491228	1
Song 2d	0.888889	0.746032	0.47619	0.492063	0	0.603175	0.52381	1
Song 2e	0.95	0.75	0.483333	0.533333	0.603175	0	0.483333	1
Song 2f	0.947368	0.719298	0.508772	0.491228	0.52381	0.483333	0	1
Song 3	1	1	1	1	1	1	1	0

<b>LUMPER</b>	Song 1	Song 2a	Song 2b	Song 2c	Song 2d	Song 2e	Song 2f	Song 3
Song 1	0	0.714286	0.823529	0.8	0.666667	0.85	0.842105	1
Song 2a	0.714286	0	0.529412	0.466667	0.619048	0.6	0.578947	1
Song 2b	0.823529	0.529412	0	0.117647	0.190476	0.15	0.105263	1
Song 2c	0.8	0.466667	0.117647	0	0.285714	0.25	0.210526	1
Song 2d	0.666667	0.619048	0.190476	0.285714	0	0.190476	0.190476	1
Song 2e	0.85	0.6	0.15	0.25	0.190476	0	0.05	1
Song 2f	0.842105	0.578947	0.105263	0.210526	0.190476	0.05	0	1
Song 3	1	1	1	1	1	1	1	0



**Figure 10:** Distances between theme strings



## Discussion

### Review of the methods used

Some of the methods used proved to be better than others. Categorization of song elements was carried out both by human and computer. However, as all computer analyses relied on previous human classification, no method used was completely objective; all of them had at least some degree of subjectivity which couldn't be avoided with just one human observer.

### Unit classification

There were some difficulties regarding categorisation of units into different types. Units from the Australian song didn't change much, but units from Atlantic changed throughout the songs. Allocating different types to these evolving units was challenging and in some cases there was just no clear border at which a new unit type replaced the old one. In these cases there was no criterion but my own decision, which depended mostly on the position of units within the song. To test if automated measures can explain human classification of units, a principal component analysis was conducted. It turned out that these measurements couldn't explain all the variation between different unit types. The measurements relied on frequency and time properties of the units, but they didn't consider the position of units within the song, or finer scaled frequency structure beyond simple energy distribution. Until such automated analysis which takes the position of units into account is invented, a human classifier is needed to account for this important unit characteristic.

### Delineation of phrases and themes

There is still no agreed method in the literature for delineation of phrases and themes. A review of the humpback whale song hierarchical structure and its classification issues does give some very useful instructions (Cholewiak *et al.* 2013), which were followed in this study whenever possible. Unfortunately, there were several situations for which these guidelines were not applicable, especially when it came to phrase delineation. For example, Cholewiak *et al.* (2013) recommends that 'phrases should be delineated in a way that minimizes the occurrence of an incomplete phrase at the end of a sequence of similar phrases', but in the Dominica song these 'hanging' phrases could not be avoided; whichever way you divide the song, there would still be a 'hanging' phrase almost at the end of every theme. The biggest difficulty I encountered was phrases that appeared to be 'transitional' (phrases that combine units from two subsequent phrases). They contained units from two phrases which were not consecutive. In these cases, a decision had to be made about when to name a new phrase type,

and it was based on a subjective interpretation. However, Cholewiak *et al.* (2013) did point out that examination of songs from multiple individuals and years is crucial for appropriately delineating phrases, so a bigger data set would probably elucidate some of the uncertainties. The same applies for themes, although they were slightly easier to deal with. Payne & McVay (1971) defined theme as sequence of similar phrases, so as Cholewiak *et al.* (2013) suggested, a new theme type was assigned every time there was a new phrase type. This means that delineation of themes heavily relied on phrase delineation.

### **Levenshtein distance method**

Levenshtein distance analysis turned out to be an effective method for comparing humpback whale songs. The beauty of it lies in its conceptual simplicity, yet it gives a clear representation of the level of similarity between different strings, whether comparing unit, phrase or theme strings. In my opinion, it is the most appropriate method for quantifying similarity (or dissimilarity) between any given song sequences. It also proved to be a very robust qualitative measure, regardless of the approach taken ('splitting' or 'lumping'). Although the LD values were higher for the 'splitter' than for the 'lumper' analysis, the clustering showed that the overall pattern of similarity between songs was the same.

### **Comparison with the previous studies**

Although there were a lot of difficulties regarding different approaches and methods used, and also some sub-optimal technical aspects, such as discrepancies in quality of recordings (different sampling rates and signal-to-noise ratios) and small sample size, the results were in accordance with the previous findings (Winn *et al.* 1981, Payne & Guinee 1983, Helweg *et al.* 1998, Cerchio *et al.* 2001, Darling & Sousa-Lima 2005) and they confirmed the hypothesis.

Even though two humpbacks from the North Atlantic were recorded in quite distant areas, they shared a relatively large portion of their song material. This conforms well with expectations derived from the literature (Winn *et al.* 1981, Guinee *et al.* 1983, Payne *et al.* 1983, Helweg *et al.* 1998, Cerchio *et al.* 2001) and suggests that these two individuals had to be in acoustic contact at some point. Perhaps they went to the same breeding or feeding grounds, or they might have shared a part of their migration routes. This study highlights the importance of horizontal cultural transmission for social and vocal learning in humpback whales.

On the other hand, it is quite hard to imagine a whale from Australia ever being in contact with whales from North Atlantic, so the opportunity for them to learn songs from each other is not really possible. Obtained results support this idea, and they clearly show that there is no similarity between any song sequences that come from these separated populations. There were a couple of shared unit types, but that was expected, as humpbacks do have a common repertoire of units (Payne & Guinee 1983).

The results of the Levenshtein distance analysis indicated that there was a substantial variation between songs of the same whale from Georges Bank at all levels (unit, phrase and theme), suggesting that, along with the differences between individuals, there is also a notable intra-individual variability.

### **Future research**

Why do humpback whales sing and why and how do these songs change? The answers to these questions are still unknown, but looking at the individual level might be the way to answer them. Instead of focusing on the big changes across populations, it would be interesting to compare more songs from whales within the same population and see how much do they differ. It would be exciting to find out is there significance in variation between individual singers, both for a group of whales and individuals themselves. One way to explore the importance of individuality would be to investigate if individual male fitness increases with more diverse and complex vocal displays.

## Conclusions

The overall conclusion is that humpbacks from North Atlantic share some of their song material and sequences, but, with the exception of two unit types, they have no similarities in their song displays with the whale from Australia. These results support the leading current theory of horizontal cultural transmission of humpback whale song.

Depending on the method and approach that was used, several conclusions can be drawn. If taking a ‘lumping’ approach and just looking at numbers of different theme, phrase and unit types, songs from the same song session seem identical, and they are similar to the other song from North Atlantic. If looking at sequences, there is a much greater variability, both between the songs of the same individual and songs from different whales. The Levenshtein distance method proved to be an excellent tool for comparing sequences. Although the overall pattern is the same, there are differences regarding the approach used. ‘Splitting’ gives much higher values of dissimilarity, which suggest a lot more variability, than ‘lumping’ approach, but these two approaches do not differ in terms of qualitative conclusions.

The principal component analysis is not able to explain all of the variation between different unit types, because it cannot take into account the position of units in the song and fine-scale frequency variation; therefore a human observer is a better choice for unit classification. It is still not perfect because, as with delineating phrases and themes, there is certainly a degree of subjectivity which can’t be avoided unless a more intensive protocol with multiple observers is used.

Hopefully in the future a more comprehensive and unambiguous set of rules for song delineation will be established that could be applied even on small data sets. Also, a much better automated method for unit categorisation is needed, which would include unit position as an important variable, together with robust frequency and duration properties of units.

## References

- Calupca, T. A., Fristrup, K. M. and Clark, C. W. "A compact digital recording system for autonomous bioacoustic monitoring." *Journal of the Acoustical Society of America* 108 (2000): 2582–2582.
- Cato, D. H. "Songs of Humpback Whales: The Australian Perspective." *Memoirs of the Queensland Museum* 30 (1991): 277–290.
- Cerchio, S., Jacobsen, J. K. and Norris, T. F. "Temporal and geographical variation in songs of humpback whales, *Megaptera novaeangliae*: Synchronous change in Hawaiian and Mexican breeding assemblages." *Animal Behaviour* 62 (2001): 313–329.
- Cholewiak, D. Evaluating the role of song in the humpback whale (*Megaptera novaeangliae*) breeding system with respect to intra-sexual interactions. 2008. Ph.D. thesis, Cornell University, Ithaca, NY.
- Cholewiak, D. M., Sousa-Lima, R. S. and Cerchio, S. "Humpback whale song hierarchical structure: Historical context and discussion of current classification issues." *Marine Mammal Science* 29 (2013): E312–E332.
- Clapham, P. J. and Mattila, D. K. "Humpback whale songs as indicators of migration routes." *Marine Mammal Science* 6 (1990): 155–160.
- Clapham, P. J. "The social and reproductive biology of humpback whales: an ecological perspective." *Mammal Review* 26 (1996): 27–49.
- Clapham, P. *Winged Leviathan: The Story of the Humpback Whale*. Colin Baxter Photography Ltd, 2013.
- Clapham, P.J., Baraff, L.S., Carlson, C.A., Christian, M.A., Mattila, D.K., Mayo, C.A., Murphy, M.A. and Pittman, S. "Seasonal occurrence and annual return of humpback whales, *Megaptera novaeangliae*, in the southern Gulf of Maine." *Canadian Journal of Zoology* 71 (1993): 440-443.
- Clark, C. W. J. and Clapham, P. "Acoustic monitoring on a humpback whale (*Megaptera novaeangliae*) feeding ground shows continual singing into late spring." *Proceedings: Biological Sciences* 271 (2004): 1051-1057.

- Darling, J. D. and Sousa-Lima, R. S. “Songs indicate interaction between Humpback whale (*Megaptera novaeangliae*) populations in the western and eastern South Atlantic Ocean.” *Marine Mammal Science* 21 (2005): 557–566.
- Darling, J. D., Gibson, K. M. and Silber, G. K. “Observations on the abundance and behavior of humpback whales (*Megaptera novaeangliae*) off West Maui, Hawaii.” Payne, R. *Communication and Behavior of Whales*. AAAS Selected Symposium 76. Westview Press, Boulder, CO., 1983. 201–222.
- Darling, J. D., Nicklin, M. E. and Nicklin, C. P. “Humpback whale songs: Do they organize males during the breeding season?” *Behaviour* 143 (2006): 1051–1101.
- Deecke, V. B. and Janik, V. M. “Automated categorization of bioacoustic signals: Avoiding perceptual pitfalls.” *Journal of Acoustical Society of America* 119 (2006): 645–653.
- Dunlop, R. A., Noad, M.J., Cato, D.H. and Stokes, D. “The social vocalization repertoire of east Australian migrating humpback whales (*Megaptera novaeangliae*).” *The Journal of the Acoustical Society of America* 122 (2007): 2893–2905.
- Eriksen, N., Miller, L. A., Tougaard, J. and Helweg, D. A. “Cultural change in the songs of humpback whales (*Megaptera novaeangliae*) from Tonga.” *Behaviour* 142 (2005): 305-328.
- Frumhoff, P. “Aberrant songs of humpback whales (*Megaptera novaeangliae*): Clues to the structure of humpback songs.” Payne, R. *Communication and behavior of whales*. AAAS Selected Symposium 76. Westview Press, Boulder, CO., 1983. 81–127.
- Garland, E. C., Gedamke, J., Rekdahl, M. L., Noad, M. J., Garrigue, C. and Gales, N. “Humpback Whale Song on the Southern Ocean Feeding Grounds: Implications for Cultural Transmission.” *PLoS ONE* 8 (2013): e79422.
- Garland, E. C., Goldizen, A. W., Rekdahl, M. L., Constantine, R., Garrigue, C., Daeschler Hauser, N., Poole, M. M., Robbins, J. and Noad, M. J. “Dynamic horizontal cultural transmission of humpback whale song at the ocean basin scale.” *Current Biology* 21 (2011): 687–691.
- Garland, E. C., Lilley, M. S., Goldizen, A. W., Rekdahl, M. L., Garrigue, C. and Noad, M. J. “Improved versions of the Levenshtein distance method for comparing sequence

- information in animals' vocalisations: tests using humpback whale song." *Behaviour* 149 (2012): 1413–1441.
- Glockner, D. A. "Determining the sex of humpback whales (*Megaptera novaeangliae*) in their natural environment." Payne, R. *Communication and behavior of whales*. AAAS Selected Symposium 76. Westview Press, Boulder, CO., 1983. 447–464.
- Green, S. R., E. Mercado III, Pack, A. and Herman, L. M. "Recurring patterns in the songs of humpback whales (*Megaptera novaeangliae*)." *Behavioural Processes* 86 (2011): 284–294.
- Guinee, L. N., Chu, K. and Dorsey, E. M. "Changes over time in the songs of known individual humpback whales (*Megaptera novaeangliae*)." Payne, R. *Communication and behavior of whales*. AAAS Selected Symposium 76. Westview Press, Boulder, CO., 1983. 59–80.
- Helweg, D.A., Cato, D.H., Jenkins, P.F., Garrigue, C. and McCauley, R.D. "Geographic variation in South Pacific humpback whale songs." *Behaviour* 135 (1998): 1–27.
- Janik, V. M. "Pitfalls in the categorization of behaviour: a comparison of dolphin whistle classification methods." *Animal Behaviour* 57 (1999): 133–143.
- Kennedy, A. S., Zerbini, A. N., Vásquez, O. V., Gandilhon, N., Clapham, P. J. and Adam, O. "Local and migratory movements of humpback whales (*Megaptera novaeangliae*) satellite-tracked in the North Atlantic Ocean." *Canadian Journal of Zoology* 92 (2014): 8–17.
- Maeda, H., Koido, T. and Takemura, A. "Principal component analysis of song units produced by humpback whales (*Megaptera novaeangliae*) in the Ryukyu region of Japan." *Aquatic Mammals* 26.3 (2000): 202–211.
- Mattila, D. K., Guinee, L. N. and Mayo, C. A. "Humpback whale songs on a North Atlantic feeding ground." *Journal of Mammalogy* 68 (1987): 880-883.
- McSweeney, D. J., Chu, K. C., Dolphin, W. F. and Guinee, L. N. "North Pacific humpback whale songs: A comparison on southeast Alaskan feeding ground songs with Hawaiian wintering ground songs." *Marine Mammal Science* 5 (1989): 139–148.

- Noad, J. M. and Cato, D. H. “A combined acoustic and visual survey of humpback whales off southeast Queensland.” *47* (2001): 507-523.
- Noad, M. J., Cato, D. H., Bryden, M. M., Jenner, M. N. and Jenner, K. C. “Cultural revolution in whale songs.” *Nature* 408 (2000): 537.
- Pace, F., Benard, F., Glotin, H., Adam, O. and White, P. “Subunit definition and analysis for humpback whale call classification.” *Applied Acoustics* 71 (2010): 1107–1112.
- Palsbøll, P.J., Clapham, P.J., Mattila, D.K., Larsen, F., Sears, R., Siegismund, H.R., Sigurjónsson, J., Vasquez, O. and Arctander, P. “Distribution of mtDNA haplotypes in North Atlantic humpback whales: the influence of behaviour on population structure.” *Marine Ecology Progress Series* 116 (1995): 1-10.
- Payne, K. and Payne, R. “Large scale changes over 19 years in songs of humpback whales in Bermuda.” *Zeitschrift für Tierpsychologie* 68 (1985): 89–114.
- Payne, K., Tyack, P. and Payne, R. S. “Progressive changes in the song of humpback whales songs (*Megaptera novaengliae*): A detailed analysis of two seasons in Hawaii.” Payne, R. *Communication and behavior of whales*. AAAS Selected Symposium 76. Westview Press, Boulder, CO., 1983. 9–57.
- Payne, R. and Guinee, L. N. “Humpbacks Whale (*Megaptera novaeangliae*) Songs as an Indicator of “Stocks”.” Payne, R. *Communication and Behavior of Whales*. AAAS Selected Symposium 76. Westview Press, Boulder, CO., 1983. 333–358.
- Payne, R. S. and McVay, S. “Songs of Humpback Whales.” *Science* 173 (1971): 585–597.
- Rendell, L. and Whitehead, H. “Culture in whales and dolphins.” *Behavioral and brain sciences* 24 (2001): 309–382.
- Rendell, L. E. and Whitehead, H. “Comparing repertoires of sperm whale codas: A multiple methods approach.” *Bioacoustics-The International Journal of Animal Sound and its Recording* 14 (2003): 61-81.
- Schulz, T. M., Whitehead, H., Gero, S. and Rendell, L. “Individual vocal production in a sperm whale (*Physeter macrocephalus*) social unit.” *Marine Mammal Science* 27 (2011): 149–166.



- Smith, J. N., Goldizen, A. W., Dunlop, R. A. and Noad, M. J. “Songs of male humpback whales, *Megaptera novaeangliae*, are involved in intersexual interactions.” *Animal Behaviour* 76 (2008): 467–477.
- Sousa-Lima, R., Norris, T., Oswald, J. and Fernandes, D. “A Review and Inventory of Fixed Autonomous Recorders for Passive Acoustic Monitoring of Marine Mammals.” *Aquatic Mammals* 39 (2003): 23-53.
- Stevick, P., Allen, J., Clapham, P.J., Katona, S.K., Larsen, F., Lien, J., Mattila, D.K., Palsbøll, P.J., Sears, R. and Sigurjónsson. “Population spatial structuring on the feeding grounds in North Atlantic humpback whales (*Megaptera novaeangliae*).” *Journal of Zoology* 270 (2006): 244–255.
- Suzuki, R., Buck, J. R. and Tyack, P. L. “Information entropy of humpback whale songs.” *The Journal of the Acoustical Society of America* 119 (2006): 1849–1866.
- Tyack, P. “Differential response of humpback whales, *Megaptera novaeangliae*, to playback of song or social sounds.” *Behavioral Ecology and Sociobiology* 13 (1983): 49-55.
- Tyack, P. “Interactions between singing Hawaiian humpback whales and conspecifics nearby.” *Behavioral Ecology and Sociobiology* 8 (1981): 105–116.
- Winn, H. E. and Winn, L. K. “The song of the humpback whale (*Megaptera novaeangliae*) in the West Indies.” *Marine Biology* 47 (1978): 97–114.
- Winn, H. E., Beamish, P., and Perkins, P. J. “Sounds of two entrapped humpback whales (*Megaptera novaeangliae*) in Newfoundland.” *Marine Biology* 55 (1979): 151–155.
- Winn, H. E., Thompson, T. J., Cummings, W. C., Hain, J., Hudnall, J., Hays, H. and Steiner, W. W. “Song of the Humpback Whale - Population Comparisons.” *Behavioral Ecology and Sociobiology* 8 (1981): 41-46.


# Appendices

**Appendix I:** Unit types


**Appendix II:** Song sequences represented by unit codes

# Appendix I


## Unit types



**a**      **b**                      **c**                      **d**                      **e**                      **f**                      **g**      **h**




**i**                      **j**                      **k**                      **l**      **m**                      **n**



**o**

**p**

**q**



**r**

**s**

**t**



